FOOD ALLERGY & ANAPHYLAXIS EMERGENCY CARE PLAN

FARE
Food Allergy Research & Education

PLACE PICTURE HERE

Name: _____ D.O.B.: _____

Allergic to: _____

Weight: _____ lbs. Asthma: ☐ **Yes (higher risk for a severe reaction)** ☐ **No**

NOTE: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. USE EPINEPHRINE.

☐ **Special Situation/Circumstance - If this box is checked, the child has an extremely severe allergy to the following food(s) _____.**

Even if the child has MILD symptoms after eating (ingesting) this food(s), Give Epinephrine immediately.

For **ANY** of the following **SEVERE** SYMPTOMS

LUNG
Shortness of breath, wheezing, repetitive cough

HEART
Pale or bluish skin, faintness, weak pulse, dizziness

THROAT
Tight or hoarse throat, trouble breathing or swallowing

MOUTH
Significant swelling of the tongue or lips

SKIN
Many hives over body, widespread redness

GUT
Repetitive vomiting, severe diarrhea

OTHER
Feeling something bad is about to happen, anxiety, confusion

OR A COMBINATION
of symptoms from different body areas

1. **INJECT EPINEPHRINE IMMEDIATELY.**
2. **Call 911.** Tell emergency dispatcher the person is having anaphylaxis and may need epinephrine when emergency responders arrive.

- Consider giving additional medications following epinephrine:
  - » Antihistamine
  - » Inhaler (bronchodilator) if wheezing
- Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
- If symptoms do not improve, or symptoms return, more doses of epinephrine can be given about 5 minutes or more after the last dose.
- Alert emergency contacts.
- Transport patient to ER, even if symptoms resolve. Patient should remain in ER for at least 4 hours because symptoms may return

MILD SYMPTOMS

NOSE
Itchy or runny nose, sneezing

MOUTH
Itchy mouth

SKIN
A few hives, mild itch

GUT
Mild nausea or discomfort

FOR **MILD SYMPTOMS** FROM **MORE THAN ONE BODY SYSTEM**, GIVE EPINEPHRINE.

FOR **MILD SYMPTOMS** FROM A **SINGLE BODY SYSTEM** (E.G. SKIN, GI, ETC.), FOLLOW THE DIRECTIONS BELOW:

1. Antihistamines may be given, if ordered by a healthcare provider.
2. Stay with the person; alert emergency contacts.
3. Watch closely for changes. If symptoms worsen, give epinephrine.

MEDICATIONS/DOSES

Epinephrine Brand or Generic: _____

Epinephrine Dose: ☐ 0.1 mg IM ☐ 0.15 mg IM ☐ 0.3 mg IM

Antihistamine Brand or Generic: _____

Antihistamine Dose: _____

Other (e.g., inhaler-bronchodilator if wheezing): _____

_____

PATIENT OR PARENT/GUARDIAN AUTHORIZATION SIGNATURE DATE

HEALTHCARE PROVIDER AUTHORIZATION SIGNATURE DATE

Form provided courtesy of **Food Allergy Research & Education** (FARE - FoodAllergy.org) - January 2023

FOOD ALLERGY & ANAPHYLAXIS EMERGENCY CARE PLAN

FARE
Food Allergy Research & Education

HOW TO USE AUVI-Q® (EPINEPHRINE INJECTION, USP), KALEO

1. Remove Auvi-Q® from the outer case. Pull off red safety guard.
2. Place black end of Auvi-Q® against the middle of the outer thigh.
3. Press firmly until you hear a click and hiss sound, and hold in place for 2 seconds.
4. Call 911 and get emergency medical help right away.



HOW TO USE EPIPEN®, EPIPEN JR® (EPINEPHRINE) AUTO-INJECTOR AND EPINEPHRINE INJECTION

1. (AUTHORIZED GENERIC OF EPIPEN®), USP AUTO-INJECTOR, MYLAN AUTO-INJECTOR, MYLAN
2. Remove the EpiPen® or EpiPen Jr® Auto-Injector from the clear carrier tube.
3. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, remove the blue safety release by pulling straight up.
4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
5. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.



HOW TO USE IMPAX EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF ADRENACLICK®), USP AUTO-INJECTOR, AMNEAL PHARMACEUTICALS

1. Remove epinephrine auto-injector from its protective carrying case.
2. Pull off both blue end caps: you will now see a red tip. Grasp the auto-injector in your fist with the red tip pointing downward.
3. Put the red tip against the middle of the outer thigh at a 90-degree angle, perpendicular to the thigh. Press down hard and hold firmly against the thigh for approximately 10 seconds.
4. Remove and massage the area for 10 seconds. Call 911 and get emergency medical help right away.



HOW TO USE TEVA'S GENERIC EPIPEN® (EPINEPHRINE INJECTION, USP) AUTO-INJECTOR, TEVA PHARMACEUTICAL INDUSTRIES

1. Quickly twist the yellow or green cap off of the auto-injector in the direction of the "twist arrow" to remove it.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, pull off the blue safety release.
3. Place the orange tip against the middle of the outer thigh at a right angle to the thigh.
4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
5. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.



HOW TO USE SYMJEPI™ (EPINEPHRINE INJECTION, USP)

1. When ready to inject, pull off cap to expose needle. Do not put finger on top of the device.
2. Hold SYMJEPI™ by finger grips only and slowly insert the needle into the thigh. SYMJEPI™ can be injected through clothing if necessary.
3. After needle is in thigh, push the plunger all the way down until it clicks and hold for 2 seconds.
4. Remove the syringe and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.
5. Once the injection has been administered, using one hand with fingers behind the needle slide safety guard over needle.

ADMINISTRATION AND SAFETY INFORMATION FOR ALL AUTO-INJECTORS:

1. Do not put your thumb, fingers or hand over the tip of the auto-injector or inject into any body part other than mid-outer thigh. In case of accidental injection, go immediately to the nearest emergency room.
2. If administering to a young child, hold their leg firmly in place before and during injection to prevent injuries.
3. Epinephrine can be injected through clothing if needed.
4. Call 911 immediately after injection.

OTHER DIRECTIONS/INFORMATION (may self-carry epinephrine, may self-administer epinephrine, etc.):

Epinephrine first, then **call 911**. Monitor the patient and call their emergency contacts right away.

EMERGENCY CONTACTS — CALL 911

RESCUE SQUAD: _____

DOCTOR: _____ PHONE: _____

PARENT/GUARDIAN: _____ PHONE: _____

OTHER EMERGENCY CONTACTS

NAME/RELATIONSHIP: _____ PHONE: _____

NAME/RELATIONSHIP: _____ PHONE: _____

NAME/RELATIONSHIP: _____ PHONE: _____

Form provided courtesy of **Food Allergy Research & Education** (FARE - FoodAllergy.org) - January 2023

WILLIAMSBURG
JAMES CITY COUNTY
PUBLIC SCHOOLS | EST. 1955

Permission to Carry and/or Self-Administer Epinephrine

Name: _____ DOB: _____

I, as the Healthcare Provider, certify that this child has a medical history of severe allergic reactions has been trained in the use of the prescribed medication(s) and is judged to be capable of carrying and self-administering this medication(s). The nurse or the appropriate school staff should be notified anytime the medication/injector is used. This child understands the hazards of sharing medications with others and has agreed to refrain from this practice.

☐ Self-Carry ☐ Self-Administer

_____	_____	_____
Healthcare Provider Signature	Print Healthcare Provider Name	Date

---

In accordance with the Code of Virginia Section 22.1-274, I agree to the following:

I will not hold the school board or any of its employees liable for any negative outcome resulting from the self-administration of said emergency medication by the student.

I understand that the school, after consultation with the parent(s) may impose reasonable limitations or restrictions upon a student's possession and/or self-administration of said emergency medication relative to the age and maturity of the student or other relevant consideration.

I understand that the school may withdraw permission to possess and self-administer the said emergency medication at any point during the school year if it is determined the student has abused the privilege of possession and self-administration or that the student is not safely and effectively self-administering the medication.

_____	_____
Parent/Guardian Signature	Date

_____	_____
Student Signature	Date